# Cold Exposure, Brown Adipose Tissue Activation, and Visceral Adiposity:

*Review of literature for book chapter VAT-TRAP © 2025 E.Leatham*

**Author:**
**Dr Ed Leatham**
**Surrey Cardiovascular Clinic**

---

# Executive Summary

For most of human history, the human thermogenic system — particularly **brown adipose tissue (BAT)** — was activated daily through environmental cold exposure. BAT is a mitochondria-rich, calorie-burning organ whose activation increases resting metabolic rate, improves insulin sensitivity, and reduces visceral adipose tissue (VAT).

Modern, temperature-controlled lifestyles suppress BAT activation, reduce D2-mediated conversion of T4→T3 within BAT, and promote VAT accumulation and metabolic inflexibility.

Multiple human studies show that **mild daily cold exposure** (e.g., **17°C for 2 hours/day, 6 weeks**) produces:

- ↑ BAT activity
- ↑ resting metabolic rate
- ↑ insulin sensitivity
- ↑ glucose oxidation
- ↓ VAT accumulation

These benefits are reversible when ambient temperatures rise.

BAT activation can also be stimulated by certain foods, notably **capsaicin** and **capsinoids** (non-pungent compounds found naturally in sweet pepper varieties), offering a potential dietary adjunct to cold-based thermogenic strategies.

This short unreviewed paper consolidates the mechanistic, clinical, and translational evidence supporting cold exposure — and selected dietary compounds — as tools for VAT reduction and metabolic health.

---

# 1. Introduction: Brown Fat and Metabolic Health

**Brown adipose tissue (BAT)** is present in adult humans and plays a critical role in:

- Non-shivering thermogenesis
- Glucose disposal
- Fatty-acid oxidation
- Insulin sensitivity
- Maintenance of metabolic flexibility

By contrast, **visceral adipose tissue (VAT)**:

- Stores calories
- Releases inflammatory cytokines
- Drives insulin resistance
- Increases cardiovascular risk

Stimulating BAT counteracts the metabolic burden imposed by VAT.

---

# 2. Mechanism: How Cold Activates BAT

Cold exposure initiates a tightly coordinated physiological cascade:

1. **Sympathetic activation**
   → noradrenaline release

2. **D2 enzyme activation**
   → increased local conversion of T4 → **T3**

3. **Rise in intracellular T3 in BAT**
   → mitochondrial uncoupling (UCP1-mediated)

4. **Heat generation**
   → increased resting metabolic rate

This mechanism increases glucose uptake, improves insulin sensitivity, and stimulates fat oxidation — all of which oppose VAT accumulation.

---

# 3. Modern Thermal Environment: A Metabolic Mismatch

Contemporary indoor environments (21–23°C year-round) suppress cold-induced thermogenesis by:

- Inhibiting D2 activity
- Reducing intracellular T3 activation in BAT
- Maintaining minimal sympathetic stimulation
- Lowering basal metabolic rate
- Supporting VAT deposition

Humans are no longer routinely exposed to the cold stimulus required to recruit and maintain BAT activity.

---

# 4. Human Evidence Supporting Cold Exposure

## 4.1 Cold-Induced BAT Recruitment

**Yoneshiro et al., J Clin Invest 2013**

Daily exposure to **17°C for 2 hours/day for 6 weeks** resulted in:

- ↑ BAT volume and metabolic activity (PET-CT)
- ↑ thermogenic capacity
- ↑ resting energy expenditure
- Greater weight reduction in BAT-positive individuals

---

## 4.2 Improved Insulin Sensitivity

**Lee et al., Diabetes 2014**

Cold acclimation:

- ↑ whole-body insulin sensitivity
- ↑ glucose disposal in BAT and muscle
- ↓ fasting insulin
- ↓ post-prandial glucose

---

## 4.3 Enhanced Non-Shivering Thermogenesis

**van der Lans et al., J Clin Invest 2013**

Just **10 days** of mild cold exposure increased:

- BAT activity
- Non-shivering thermogenesis
- Resting metabolic rate

---

## 4.4 Cold Exposure for Type 2 Diabetes Prevention

**Schrauwen & van Marken Lichtenbelt, Diabetologia 2016**

This review establishes that chronic mild cold exposure improves glycaemic control and may be a **physiological adjunct therapy** for preventing and managing type 2 diabetes.(1)

---

# 5. Dietary BAT Activators (Capsaicin and Capsinoids)

Beyond cold exposure, certain dietary compounds can stimulate BAT activation.
The most extensively studied are **capsaicin** (from chilli peppers) and **capsinoids**, non-pungent analogues found in certain sweet pepper varieties.

**Mechanism:**

- Capsaicin and capsinoids activate **TRPV1 receptors**
- This increases sympathetic outflow to BAT
- D2 activity rises
- Intracellular T3 production increases
- BAT thermogenesis increases
- Resting metabolic rate increases

**Key Human Study:**
**Yoneshiro et al., AJCN 2012 (capsinoid trial)** (2)

- Capsinoid ingestion increased BAT metabolic activity
- PET-CT confirmed increased ^18F-FDG uptake
- Energy expenditure increased
- Fat oxidation increased
- Effects were seen even without cold exposure

This demonstrates that **capsaicin and capsinoids can mimic mild cold exposure**, making them a valuable dietary adjunct to improve metabolic rate and reduce VAT accumulation.

---

# 6. Why Cold Exposure Reduces VAT

## 6.1 Increased Caloric Expenditure

Non-shivering thermogenesis increases baseline energy use.

## 6.2 Improved Insulin Sensitivity

Reducing hyperinsulinaemia reduces VAT deposition.

## 6.3 Enhanced Glucose Uptake

BAT acts as a glucose sink.

## 6.4 Increased Fat Oxidation

BAT oxidises fatty acids rapidly during thermogenesis.

### 6.5 Restoration of D2–T3 Signalling

Obesity reduces D2 activity → low intracellular T3
Cold exposure increases D2 → restores T3 signalling

---

# 7. Practical Clinical Recommendations

### 7.1 Lower Indoor Temperatures

- Aim for **18–19°C** during the day
- **16–18°C** at night

### 7.2 Light Cold Exposure

- Walk outside with one fewer layer
- Avoid overheating indoors
- Gradual acclimatisation recommended

### 7.3 Optional Cold-Enhanced Strategies

- Cool showers
- Cooling vests
- Intermittent cold periods

### 7.4 Complementary Lifestyle Measures

Cold exposure synergises with:

- Resistance training
- Protein-intensive diets
- Glycaemic control (CGM-guided)
- VAT monitoring (waist–height ratio)

---

# 8. Public Health Perspective

Cold exposure is:

- Safe
- Free
- Scalable
- Environmentally beneficial (lower heating use)
- Effective in improving metabolic health

It represents an underutilised metabolic tool with potential population-level benefits for reducing VAT, obesity, insulin resistance, and diabetes.

---

# 9. Conclusion

BAT is a powerful metabolic organ capable of improving insulin sensitivity, increasing metabolic rate, and reducing VAT. Modern living suppresses BAT activation, but **mild daily cold exposure** and certain dietary compounds (capsaicin, capsinoids) can safely restore this ancient metabolic pathway.

Turning the thermostat down is a **simple, evidence-based, physiologically grounded** intervention to improve metabolic health, reduce VAT, and support cardiometabolic prevention.

---

# References


### 1. Yoneshiro T, Aita S, Matsushita M, et al.

Recruited brown adipose tissue as an antiobesity agent in humans.
*J Clin Invest.* 2013;123(8):3404–3408.
DOI: 10.1172/JCI67803
https://pubmed.ncbi.nlm.nih.gov/23867622/
https://www.jci.org/articles/view/67803

YouTube video explainer

### 2. Lee P, Smith S, Linderman J, et al.

Temperature-acclimated brown adipose tissue modulates insulin sensitivity in humans.
*Diabetes.* 2014;63(11):3686–3698.
DOI: 10.2337/db14-0513
https://pubmed.ncbi.nlm.nih.gov/24954193/
https://pmc.ncbi.nlm.nih.gov/articles/PMC4207391/

### 3. van der Lans AAJJ, Hoeks J, Brans B, et al.

Cold acclimation recruits human brown fat and increases nonshivering thermogenesis.
*J Clin Invest.* 2013;123(8):3395–3403.
DOI: 10.1172/JCI68993
https://pubmed.ncbi.nlm.nih.gov/23867626/
https://pmc.ncbi.nlm.nih.gov/articles/PMC3726172/

### 4. Schrauwen P, van Marken Lichtenbelt WD.

Combatting type 2 diabetes by turning up the heat.
*Diabetologia.* 2016;59(11):2269–2279.
DOI: 10.1007/s00125-016-4068-3
https://pubmed.ncbi.nlm.nih.gov/27591854/
https://pmc.ncbi.nlm.nih.gov/articles/PMC5506100/

### 5. Yoneshiro T, Aita S, Kawai Y, et al.

Nonpungent capsaicin analogs (capsinoids) increase energy expenditure through activation of brown adipose tissue in humans.
*Am J Clin Nutr.* 2012;95(4):845–850.
DOI: 10.3945/ajcn.111.022921
https://pubmed.ncbi.nlm.nih.gov/22378725/
https://academic.oup.com/ajcn/article/95/4/845/4576742

**6. Cannon B, Nedergaard J.**

Brown adipose tissue: function and physiological significance.
*Physiol Rev.* 2004;84(1):277–359.
DOI: 10.1152/physrev.00015.2003
https://pubmed.ncbi.nlm.nih.gov/14715917/
https://journals.physiology.org/doi/10.1152/physrev.00015.2003

**7. Silva JE.**

Thermogenic mechanisms and their hormonal regulation.
*Physiol Rev.* 2006;86(2):435–464.
DOI: 10.1152/physrev.00009.2005
https://pubmed.ncbi.nlm.nih.gov/16601266/
https://journals.physiology.org/doi/10.1152/physrev.00009.2005

**8. Virtue S, Vidal-Puig A.**

Adipose tissue expandability, lipotoxicity and metabolic syndrome – an allostatic perspective.
*Biochim Biophys Acta.* 2010;1801(3):338–349.
DOI: 10.1016/j.bbalip.2009.12.006
https://pubmed.ncbi.nlm.nih.gov/20056169/

**9. Wu H, Ballantyne CM.**

Metabolic inflammation and insulin resistance in obesity.
*Circ Res.* 2020;126(11):1549–1564.
DOI: 10.1161/CIRCRESAHA.119.315896
https://pubmed.ncbi.nlm.nih.gov/32437299/
https://pmc.ncbi.nlm.nih.gov/articles/PMC7250139/